



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C07K 1/00, C07K 14/00,</b> <b>C07K 17/00, C12N 9/00,</b> <b>C12N 9/12, C12P 21/06,</b> <b>C30B 7/00, C30B 21/02,</b> <b>C30B 28/06, G01N 31/00,</b> <b>G01N 33/00, G01N 33/53,</b> <b>G01N 33/567</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/14105</b> <b>(43) International Publication Date:</b> <b>16 March 2000 (16.03.2000)</b>
<b>(21) International Application Number:</b> PCT/US99/20319 <b>(22) International Filing Date:</b> 07 September 1999 (07.09.1999) <b>(30) Priority Data:</b> 60/099,503 08 September 1998 (08.09.1998) US <b>(60) Parent Application or Grant</b> AGOURON PHARMACEUTICALS, INC. [/]; (). MCTIGUE, Michele, A. [/]; (). PINKO, Chris [/]; (). PARAST, Camran, V. [/]; (). GEHRING, Michael, R. [/]; (). KAN, Chen-Chen [/]; (). APPELT, Krzysztof [/]; (). WICKERSHAM, John, A. [/]; (). SHOWALTER, Richard, Edward [/]; (). TEMPCYZK-RUSSELL, Anna-Maria [/]; (). MROCZKOWSKI, Barbara [/]; (). VILLAFRANCA, Jesus, Ernesto [/]; (). HERBERT, Toni-Junell; ().		<b>Published</b>
<b>(54) Title: MODIFICATIONS OF THE VEGF RECEPTOR-2 PROTEIN AND METHODS OF USE</b> <b>(54) Titre: MODIFICATION DE LA PROTEINE RECEPTEUR 2 DE VEGF ET SES METHODES D'UTILISATION</b>  <b>(57) Abstract</b> <p>A 2.4 Å crystal structure of a protein construct containing the catalytic kinase domain of vascular endothelial growth factor receptor 2 (VEGFR2/KDR), a key enzyme in angiogenesis, has been determined in an unliganded, phosphorylated state. This protein construct, contains a modified catalytic linker and has comparable in vitro kinase activity to constructs containing the entire KID. The resulting construct retains comparable in vitro kinase activity to that of the wild-type KID, and more importantly, allows complete crystallization of the protein such that it may be characterized by X-ray crystallography. The present invention further discloses the use of x-ray crystallographic data for identification and construction of possible therapeutic compounds in the treatment of various disease conditions.</p> <b>(57) Abrégé</b> <p>On a identifié une structure cristalline de 2,4 Å d'une construction protéique contenant le domaine catalytique de la kinase, du récepteur 2 du facteur de croissance de l'endothélium vasculaire (VEGFR2/KDR), une enzyme clé dans l'angiogénèse, à l'état phosphorylé et non ligand. Cette construction protéique contient un segment de liaison catalytique et possède une activité de kinase in vitro comparable aux constructions contenant le domaine d'insertion de la kinase (KID). La construction résultante conserve une activité de kinase in vitro comparable à celle du KID du type sauvage, et mieux encore, permet la cristallisation complète de la protéine de sorte qu'elle puisse être caractérisée par cristallographie par rayons X. L'invention porte encore sur l'utilisation de données de cristallographie par rayons X, pour l'identification et la construction d'éventuels composés thérapeutiques dans le traitement de divers états pathologiques.</p>		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : C07K 1/00, 14/00, 17/00, C12N 9/00, 9/12, C12P 21/06, C30B 7/00, 21/02, 28/06, G01N 31/00, 33/00, 33/53, 33/567</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 00/14105</b>  (43) International Publication Date: 16 March 2000 (16.03.00)</p>
<p>(21) International Application Number: PCT/US99/20319  (22) International Filing Date: 7 September 1999 (07.09.99)  (30) Priority Data: 60/099,503 8 September 1998 (08.09.98) US  (71) Applicant: AGOURON PHARMACEUTICALS, INC. [US/US]; Legal Dept., 10350 North Torrey Pines Road, La Jolla, CA 92037 (US).  (72) Inventors: MCTIGUE, Michele, A.; 515 Hermes Avenue, San Diego, CA 92024 (US). PINKO, Chris; 7953 Playmor Terrace, San Diego, CA 92122 (US). PARAST, Cam- ran, V.; 1855 Diamond Street #5-218, San Diego, CA 92109 (US). GEHRING, Michael, R.; 5468 Collier Av- enue, San Diego, CA 92115 (US). KAN, Chen-Chen; 13951 Durango Drive, Del Mar, CA 92014 (US). APPELT, Krzysztof; 14500 Maplewood Street, Poway, CA 92064 (US). WICKHRSHAM, John, A.; 2426 Shadyridge Avenue, Escondido, CA 92029 (US). SHOWALTER, Richard, Ed- ward; 13329 Goldentop Drive, Lakeside, CA 92040 (US). TEMPCYZK-RUSSELL, Anna-Maria; 4785 Camino Bor- rego, San Diego, CA 92130 (US). MROCZKOWSKI, Bar- bara; 1066 Stratford Drive, Encinitas, CA 92024 (US). VIL-</p>	<p>LAFRANCA, Jesus, Ernesto; 1282 Upas Street, San Diego, CA 92103 (US).  (74) Agents: HERBERT, Toni-Junell; Shanks &amp; Herbert, Suite 306, TransPotomac Plaza, 1033 North Fairfax Street, Alexandria, VA 22314 (US) et al.  (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SI, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	
<p>(54) Title: MODIFICATIONS OF THE VEGF RECEPTOR-2 PROTEIN AND METHODS OF USE</p>		
<p>(57) Abstract  A 2.4 Å crystal structure of a protein construct containing the catalytic kinase domain of vascular endothelial growth factor receptor 2 (VEGFR2/KDR), a key enzyme in angiogenesis, has been determined in an unliganded, phosphorylated state. This protein construct, contains a modified catalytic linker and has comparable <i>in vitro</i> kinase activity to constructs containing the entire KID. The resulting construct retains comparable <i>in vitro</i> kinase activity to that of the wild-type KID, and more importantly, allows complete crystallization of the protein such that it may be characterized by X-ray crystallography. The present invention further discloses the use of x-ray crystallographic data for identification and construction of possible therapeutic compounds in the treatment of various disease conditions.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	RS	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**Description**

5

10

15

20

25

30

35

40

45

50

55

5  
10  
15  
20  
25  
**Modifications of The VEGF Receptor-2 Protein And Methods Of Use****TECHNICAL FIELD AND INDUSTRIAL APPLICABILITY OF INVENTION**

5 The present invention discloses the isolation of a key portion of the catalytic kinase region of vascular endothelial growth factor receptor 2 or VEGFR-2 through cloning, sequencing and x-ray crystallography. Also disclosed is the deletion of various amino acid residues from an area of the catalytic region called the kinase insert domain (KID). The resulting polypeptide retains comparable *in vitro* kinase activity to that of the wild-type KID and is not necessary for the catalytic activity of the polypeptide, and more importantly, allows complete crystallization of the protein such that it may be characterized by X-ray crystallography. The present invention further discloses x-ray crystallography data useful for identification and construction of therapeutic compounds in the treatment of various disease conditions associated with VEGFR-2.

**BACKGROUND OF THE INVENTION**

30 Many physiological events including embryogenesis, organ development, estrus, and wound healing require vascular growth and remodeling (Folkman et al., (1992) J. Biol. Chem. 267, 10931-10934; Risau, W. (1995) FASEB J. 9, 926-933.). In addition to these beneficial processes, angiogenesis is also involved in the proliferation of disease states such as tumor growth, metastasis, psoriasis, rheumatoid arthritis, macular degeneration and retinopathy (Pepper, M.S., (1996) Vasc. Med. 1, 259-266; Kuiper et al., (1998) Pharmacol. Res. 37, 1-16, 1998; Kumar and Fidler, (1998) In Vivo 18, 27-34; Szekanecz et al., (1998) J. Invest. Med. 46, 27-41; Tolentino and Adamis, (1988) Int. Ophthalmol. Clin. 38, 77-94. Of the signaling pathways known to influence vascular formation, these involving vascular endothelial growth factor (VEGF) have been shown to be essential and selective for vascular endothelial cells (Dvorak et al., (1995) Am. J. Path. 146, 1029-1039; Thomas, K., (1996) Cell 271, 603-606; Ferrara N. and Davis-Smyth, (1997) Endocrine Rev. 18, 4-25). The therapeutic potential of inhibiting the VEGF pathway has been directly demonstrated by anti-VEGF monoclonal antibodies which were active against a variety of human tumors (Borgström et al,

(1996) Cancer Res. 56, 4032-4039) and ischemic retinal disease (Adamis et al., (1996) Arch. Ophthalmol. 114, 66-71).

Normal vasculogenesis and angiogenesis play important roles in a variety of physiological processes such as embryonic development, wound healing, organ regeneration and female reproductive processes such as follicle development in the corpus luteum during ovulation and placental growth after pregnancy (Folkman & Shing, 1992). Uncontrolled vasculogenesis and/or angiogenesis has been associated with diseases, such as diabetes, as well as malignant solid tumors that rely on vascularization for growth. Klagsburn & Soker, (1993) Current Biology 3(10):699-702; Folkham, (1991) J. Natl. Cancer Inst. 82:4-6; Weidner, et al., (1991) New Engl. J. Med. 324:1-5.

Several polypeptides with *in vitro* endothelial cell growth promoting activity have been identified. Examples include acidic and basic fibroblastic growth factor (FGF), vascular endothelial growth factor (VEGF) and placental growth factor. Unlike FGF, VEGF has recently been reported to be an endothelial cell specific mitogen (Ferrara & Henzel, (1989) Biochem. Biophys. Res. Comm. 161:851-858; Vaisman et al., (1990) J. Biol. Chem. 265:19461-19566).

Thus, identification of the specific receptors to which VEGF binds is important to understanding of the regulation of endothelial cell proliferation. Two structurally related tyrosine kinases have been identified to bind VEGF with high affinity: the flt-1 receptor (Shibuya et al., (1990) Oncogene 5:519-524; De Vries et al., (1992) Science 255:989-991) and the KDR/FLK-1 receptor, discussed herein. Consequently, it had been surmised that RTKs may have a role in the modulation and regulation of endothelial cell proliferation.

Recent disclosures, such as information set forth in U.S. Patent Application Ser. Nos. 08/193,829, 08/038,596 and 07/975,750, strongly suggest that VEGF is not only responsible for endothelial cell proliferation, but also is the prime regulator of normal and pathological angiogenesis. See generally, Klagsburn & Soker, (1993) Current Biology 3:699-702; Houck, et al., (1992) J. Biol. Chem. 267:26031-26037.

5 VEGF is a homodimeric cytokine that is expressed in at least four splice-variant forms of 121-206 residues (Ferrara and Davis-Smyth, 1997). Vascular  
endothelial cells express at least two high-affinity receptors for VEGF: VEGF-R1/Flt-1  
10 and VEGFR-2/KDR. VEGF-R1 and VEGFR-2 are receptor tyrosine kinases each  
5 comprised of an extracellular domain that contains 7 immunoglobulin-like segments  
and binds VEGF, a short membrane spanning region, and a cytosolic domain  
possessing tyrosine kinase activity. The kinase domain directly follows the  
15 extracellular and juxtamembrane regions and is itself followed by another domain  
(post-kinase domain), which may function in binding of other proteins for signal  
20 transduction. These two receptors appear to have different signaling pathways and  
functions with VEGFR-2 being of primary importance in mitosis of endothelial cells  
(Waltenberger et al., (1994) J. Biol. Chem. 269, 26988-26995; Seetharm et al.,  
(1995) Oncogene 10, 135-147; Shalaby et al., (1995) Nature 376, 576-579).

25 Both FGF and VEGF are potent angiogenic factors which induce formation of  
new capillary blood vessels. Transfection of human breast carcinoma cell line MCF-7  
with FGF resulted in cell lines that form progressively growing and metastatic tumors  
when injected (s.c.) into nude mice. FGF may play a critical role in progression of  
30 breast tumors to an estrogen-independent, anti-estrogen resistant metastatic  
20 phenotype (McLeskey et al., (1993) Cancer Res. 53: 2168-2177). Breast tumor cells  
exhibited increased neovascularization, increased spontaneous metastasis and more  
rapid growth in vivo than did the non-transfected tumors. FGF has been shown to be  
35 transforming in NIH-3T3 cells and implicated in tumorigenesis and metastasis of  
mouse mammary tumors. FGF overexpression conferred a tumorigenic phenotype on  
25 a human adrenal carcinoma cell line suggesting that FGF's may also play a role in the  
transformation of epithelial cells. Polyclonal neutralizing antibodies to FGF inhibited  
40 tumor growth in Balb/c nude mice transplanted with K1000 cells (transfected with the  
leader sequence of bFGF) which form tumors in these mice (Hori et al., (1991)  
Cancer Res. 51: 6180-9184).

30  
45 Due to the role of FGF in neovascularization, tumorigenesis and metastasis,  
there is a need in the art for FGF inhibitors as potent anti-cancer agents that exert  
their anti-FGF activity by preventing intracellular signaling of FGF.

5

10

15

10

20

25

30

20

35

40

45

50

55

VEGF, by contrast, is an endothelial cell-specific mitogen and an angiogenesis inducer that is released by a variety of tumor cells and expressed in human tumor cells *in situ*. Unlike FGF, transfection of cell lines with a cDNA sequence encoding VEGF, did not promote transformation, but did facilitate tumor growth *in vivo* (Ferrara, N., and Davis-Smyth, T. (1997)). Furthermore, administration of a polyclonal antibody which neutralized VEGF also inhibited growth of human rhabdomyosarcoma, glioblastoma multiforme and leiomyosarcoma cell lines in nude mice (Kim et al., (1993) Nature 362: 841-843).

In view of the importance of receptor tyrosine kinases (RTKs) to the control, regulation and modulation of endothelial cell proliferation and potentially vasculogenesis and/or angiogenesis, many attempts have been made to identify RTK "inhibitors" using a variety of approaches, including the use of mutant ligands (U.S. Patent No. 4,966,849), soluble receptors and antibodies (Application No. WO 94/10202; Kendall & Thomas, (1994) Proc. Natl. Acad. Sci. 90:10705-09; Kim, et al., 1993), RNA ligands (Jellinek, et al., (1994) Biochemistry 3:10450-56), protein kinase C inhibitors (Schuchter, et al., (1991) Cancer Res. 51:682-687); Takano, et al., (1993) Mol. Bio. Cell 4:358A; Kinsella, et al., (1992) Exp. Cell Res. 199:56-62; Wright, et al., (1992) J. Cellular Phys. 152:448-57) and tyrosine kinase inhibitors (WO 94/03427; WO 92/21660; WO 91/15495; WO 94/14808; U.S. Pat. No. 5,330,992; Mariani, et al., (1994) Proc. Am. Assoc. Cancer Res. 35:2268).

More recently, attempts have been made to identify small molecules which act as tyrosine kinase inhibitors. For example, bis monocyclic, bicyclic or heterocyclic aryl compounds (PCT WO 92/20642), vinylene-azaindole derivatives (PCT WO 94/14808) and 1-cyclopropyl-4-pyridyl-quinolones (U.S. Pat. No. 5,330,992) have been described generally as tyrosine kinase inhibitors. Styryl compounds (U.S. Pat. No. 5,217,999), styryl-substituted pyridyl compounds (U.S. Pat. No. 5,302,606), certain quinazoline derivatives (EP Application No. 0 566 266 A1), selenoindoles and selenides (PCT WO 94/03427), tricyclic polyhydroxylic compounds (PCT WO 92/21660) and benzylphosphonic acid compounds (PCT WO 91/15495) have been

described as compounds for use as tyrosine kinase inhibitors for use in the treatment of cancer. None of these compounds, however, have been previously associated with the enzymatic function of the VEGFR-2 receptor. Likewise, none of these compounds have been associated with regulation of vasculogenesis and/or angiogenesis.

Therefore, there is a need in the art to develop small molecule antagonists of the PDGF, FGF, EGF and VEGF pathways individually or as a group. Moreover, if these cytokines signal through a common second messenger pathway within the cell, such antagonists will have broad therapeutic activity to treat or prevent the progression of a broad array of diseases, such as coronary restenosis, tumor-associated angiogenesis, atherosclerosis, autoimmune diseases, acute inflammation, certain kidney diseases associated with proliferation of glomerular or mesangial cells, and ocular diseases associated with retinal vessel proliferation. The present invention was made by discovering a common signaling mechanism, a group of active therapeutic agents, shown to be active by a large number of and variety of predictive assays, and discovering a common intracellular signaling intermediate.

Based on sequence homology and overall domain structure, VEGFRs belong to the platelet-derived growth factor receptor family (PDGFR) which also includes PDGFR, PDGFR, the stem cell growth factor receptor (c-kit), and the colony stimulating factor-1 receptor (CSF-1R/c-fms) (van der Geer et al., (1994) Ann. Rev. Cell Biol. 10, 251-337). Compared to other protein kinases, members of this family contain an insert of approximately 65-97 residues, termed the kinase insert domain (KID), within the catalytic kinase domain relative to other protein kinases. Within the PDGFR family the KIDs are of varying length and low sequence homology. Deletion or mutation of the KID from PDGFR, PDGFR, c-kit, and CSF-1R have indicated that this domain is not necessary for intrinsic kinase activity but that it is important for the binding of other proteins involved in signal transduction, via autophosphorylation of KID tyrosine residues (Taylor et al., (1989) EMBO J. 8, 2029-2037; Heidaran et al., (1991) Mol. Cell. Biol. 11, 134-142; Yu et al., (1991) Mol. Cell. Biol. 11, 3780-3785; Kaziauskas et al., (1992) Mol. Cell. Biol. 12, 2534-2544; Lev et al., (1992) Proc. Natl. Acad. Sci. USA 89, 678-682; Reedjik et al., (1992) EMBO J. 11, 1365-1372;

5 Bazenet et al., (1996) Mol. Cell. Biol. 16, 6926-6936). Although the signaling  
pathways and the specific role of the KID are still not fully determined for VEGFRs,  
the VEGFR-2 KID does contain two tyrosines which are known to be  
10 autophosphorylation sites (Dougher-Vermazen et al., (1994) Biochem. Biophys. Res.  
5 Comm. 205, 726-738).

Since the determination of the first cyclic AMP-dependent protein kinase  
15 (cAPK) structure (Knighton et al., (1991) Science 253, 407-413) a variety of protein  
kinase structures have been reported (reviewed in Johnson et al., (1996) Cell 85,  
10 149-158). Among the receptor protein tyrosine kinases (RTKs), structures of the  
kinase domain of the insulin receptor (IRK) (Hubbard, et al., (1994) Nature 372, 746-  
20 754; Hubbard, (1997) EMBO J. 16, 5572-5581) and the fibroblast growth factor  
receptor-1 (FGFR1) (Mohammadi et al., (1996) Cell 86, 577-87; Mohammadi et al.,  
(1997) Science 276, 955-960) have been determined.

#### SUMMARY OF THE INVENTION

30 The present invention discloses the generation, kinetic characterization, and  
20 structure determination of a modified kinase domain of the VEGFR-2 protein,  
containing 18 residues of the 68 residue KID. This 2.4 Å crystal structure of the  
phosphorylated VEGFR-2 catalytic domain is the first reported structure of a kinase  
35 domain of the PDGFR family. This structure provides insights into the orientation of  
the KID domain of VEGFR-2 which may be relevant to other PDGFR family members.  
25 Furthermore, as inhibition of VEGFR-2 kinase has broad clinical applications, this  
structure provides a three-dimensional description of the target for structure-based  
40 design of small molecule VEGFR-2 inhibitors as therapeutic agents.

It is an object of the present invention to disclose an effective method for  
45 30 screening candidate compounds that are specifically agonists or antagonists of  
various proteins which can be included in the receptor tyrosine kinase family (RTK) by  
crystallizing RTKs and particularly the VEGFR-2 receptor in order to use molecular

5

modeling of the x-ray crystallography data to model the binding of candidate compounds.

10

15

20

25

30

35

40

45

There is disclosed a method for designing and screening potentially  
5 therapeutic compounds with activities such as: (1) inhibiting new blood vessel  
formation that is useful for treating or preventing progression of diabetic retinopathy,  
cavernous hemangiomas, Kaposi's sarcoma, tumors composed of endothelial-like  
cells, and growth of cancer cells by preventing their development of a new blood  
supply; (2) suppressing development of kidney diseases due to cytokine induced  
10 proliferation of mesangial cells and/or glomerular epithelial cells that is useful for  
treating or preventing progression of diabetic glomerulosclerosis and other  
glomerulonephritis of various types and etiologies; (3) preventing joint destruction  
accompanying rheumatoid arthritis due to proliferation of synovial cells; (4)  
suppressing manifestations of psoriasis due to proliferation of keratinocytes and  
15 accumulation of inflammatory cells; (5) suppressing accelerated atherogenesis  
involved in restenosis of coronary vessels or other arterial vessels following  
angioplasty; (6) suppressing atherogenesis, coronary artery disease and other  
vasculopathies due to atherogenesis; and (7) suppressing tumor growth via paracrine  
or autocrine mediated responses to other cytokines such as PDGF, FGF EGF or  
20 VEGF that is useful for treating or preventing progression of tumors such as breast  
cancer stimulated through overexpression of her-2-neu receptor, wherein the  
inventive method comprises administering a compound that inhibits signal  
transduction.

25 The present invention is useful in developing methods that are used in the  
iterative drug design process. The process identifies potential agonists and  
antagonists to VEGFR-2 by *de novo* design of novel drug candidate molecules which  
bind to the VEGFR-2 receptor to improve their potency. The x-ray crystallographic  
coordinates disclosed herein, will allow generation of 3-dimensional models of the  
30 catalytic site and drug binding site of the VEGFR-2 protein.

50

*De novo* design primarily consists of the generation of molecules via the use  
of computer programs which build and link fragments or atoms into a site based upon

55

5           steric and electrostatic complementarity, without reference to substrate analog  
structures. The drug design process begins after the structure of a target RTK is  
solved to at least a resolution of 2.8 Å. Refinement of the structure to a resolution of  
10       2.5 Å or better, with "fixed" water molecules in place provides more optimal conditions  
5       to undertake drug design.

15           It is another object of this invention to identify KIDs of proteins in the RTK  
family and develop deletions in said KIDs such that the proteins will be crystallizable  
and suitable for measurement by x-ray crystallographic means.

10           It is a further object of this invention to disclose a process whereby KID  
20       regions from a member of the RTK family of genes such as PDGF, EGF, VEGF and  
others are modified by deletion of amino acids from the KID regions so as to impart  
favorable physical characteristics of the resulting polypeptide product. Examples of  
15       such favorable physical characteristics are increased solubility, greater stability to  
temperature variations making the polypeptide suitable for analysis by nuclear  
magnetic resonance, high throughput screening, biochemical characterizations, x-ray  
crystallography, calorimetry and other diagnostic means.

30           It is yet another object of this invention to developing screening methods used  
in the drug design process of potential agonists and antagonists to proteins in the  
RTK family by de novo design of novel drug candidate molecules with potentially  
35       nanomolar potencies. The x-ray crystallographic coordinates disclosed based on the  
deletion mutated KIDs and various other deletions of said proteins in the RTK family,  
25       will allow generation of 3-dimensional models of the active binding site of the proteins  
in the RTK family.

#### BRIEF DESCRIPTION OF THE DRAWINGS

45       30       Figure 1. Secondary structure assignments (as given by Procheck) for the catalytic  
domain of VEGFR2 and sequence alignment with other representative receptor  
tyrosine kinases. helices are designated as  $\alpha$ -1,  $\beta$  strands are designated as 1-8.  
The site of 50 residue deletion in VEGFR2 50 is indicated by |. The site of the E990V

mutation in VEGFR2D50 is denoted by an \*. Sequences are from: VEGFR2 (reported here); FGFR1 (Swiss protein database #P11362); IRK (EMBL protein database #A18657; numbering as in Mohammadi et al., 1996); VEGFR1 (Swiss protein database #P17948); PDGFR (Swiss protein database #P17948).

Figure 2. Overall fold of VEGFR2 50P, FGFR1, and IRKP.

Backbone representation of structures of the kinase domains of (A) VEGFR2 (VEGFR2 50P), (B) FGFR1 (molecule A of PDB entry 1FGK, Mohammadi et al., 1996), and (C) IRKP (PDB entry 1IR3, Hubbard et al., 1997). The views shown in A, B, and C are identical views generated from superpositions of the C-terminal domains. The positions of the termini are denoted by N and C. The nucleotide-binding loop (orange), kinase insert domain (pink), and activation loop (yellow) are highlighted. In (C) the bound AMP-PNP is shown in green and the peptide substrate is shown in red. Figure prepared with INSIGHT II.

Figure 3. Catalytic site of VEGFR2 50P and IRKP.

Cross section of the catalytic site of (A) VEGFR2 50P and (B) IRKP (PDB entry 1IR3; Hubbard et al., 1997) structures. Atoms are colored by element type: carbon (green), oxygen (red), nitrogen (blue), sulfur (yellow), phosphorous (pink), and magnesium ion (orange). (A) includes only protein atoms. (B) includes protein atoms, AMP-PNP atoms, and  $Mg^{2+}$  ions. Figure generated using INSIGHT II.

Figure 4. Nucleotide binding site of VEGFR2 50P and FGFR1.

Stereo view showing C trace and some sidechains of a superposition of the nucleotide binding sites of the VEGFR2 50P and the FGFR1-(AMP-PCP) complex (molecule B, Mohammadi et al., 1996) structures. The superposition was done using C positions of helices (D, E, F, G, H, and I) of the C-terminal lobes. Carbon atoms of VEGFR2 50P are shown in yellow and carbon atoms of FGFR1 are shown in purple. The coloring for other protein atoms is: oxygen (red), nitrogen (blue), and sulfur (green). The AMP-PCP in the FGFR1 structure is depicted in orange. Labels correspond to VEGFR2 50P residues. Figure created with Xfit (McRee et al., (1992) *J. Mol. Graph.* 10, 44-46.).

Figure 5. Electron density map of the kinase insert domain area of VEGFR2 50P.

Stereo view of a  $2F_o - F_c$  map computed at 2.4Å and contoured at 1.2 and superimposed with the refined model. Carbon atoms are yellow, oxygen atoms red, and nitrogen atoms are blue. Water molecules are depicted as red crosses. Figure created with Xfit (McRee et al., 1992).

Figure 6. Kinase insert domain of VEGFR2 50P. Stereo cross section showing the ordered residues of the kinase insert domain of VEGFR2 50P. Carbon atoms are yellow, oxygen atoms are red, nitrogen atoms are blue, and sulfur atoms are green. View is rotated roughly 180° from Figure 5. Figure created with Xfit (McRee et al., 1992).

Figure 7. Resulting X-ray crystallography coordinates for VEGFR-2 based on the method disclosed in the crystallization and data collection section.

#### DETAILED DESCRIPTION AND PREFERRED EMBODIMENTS OF THE INVENTION

##### Cloning of The VEGFR-2 Protein

The coding sequence (Terman et al., (1992) Biochem Biophys. Res. Commun. 187, 1579-86) for the cytoplasmic domain of the VEGFR-2 was amplified by PCR (Mullis et al., (1986). Biotechnology 24, 17-27) from a human aorta cDNA pool (Clontech Palo Alto, CA ). Two overlapping sequences were amplified independently. Vcyt (residues M806-V1356), which represented the entire cytoplasmic domain, and Vcat (residues C817M-G1191), with boundaries based upon a primary amino acid sequence alignment with the insulin receptor kinase catalytic domain (Wei et al., (1995) J. Biol. Chem. 270, 8122-8130).

The PCR oligonucleotide primer sequences for Vcyt were:

Vcyt5 5'-CAGCATATGGATCCAGATGAACTCCCATTTGG-3' (Seq. ID No. 1) and Vcyt3 5'-GCGGTCGACTTAAACAGGAGGAGAGCTCAGTGTG-3' (Seq. ID No. 2).

The PCR oligonucleotide primer sequences for the Vcat were:

Vcat5 5'-GCACATATGGAACGACTGCCTTATGATGCCAGC-3' (Seq. ID No. 3) and  
Vcat3 5'-CCTGTCGACTTATCCAGAATCCTCTCCATGCTCAAAG-5' (Seq. ID No. 4).

The amplified DNA was digested with the restriction enzymes NdeI and Sall,  
ligated into the *E coli* plasmid pET24a (Novagen Madison, WI) and sequence verified.  
When compared to the original VEGFR-2 sequence in Genbank, (Accession number  
346345) two nucleotide differences were noted that resulted in codon changes  
(Glu848-Val and Asn835-Lys) in both Vcyt and Vcat. Our sequence agrees with  
subsequent VEGFR-2 Genbank submissions (Accession numbers 2655412 and  
3132833).

Mutations were introduced by oligonucleotide site directed mutagenesis  
(Kunkel, 1985) using the Muta-Gene in vitro Mutagenesis Kit from (Bio-Rad Hercules,  
CA). The Vcat DNA fragment was subcloned from the pET24a vector using an NdeI-  
XhoI digest into the vector pMGH4 (Schoner et al., 1986, Kan et al., 1992) and this  
vector was used to generate the ssDNA uracil template (minus strand) in *E. coli* strain  
CJ236 supplied in the kit. An oligo (5'-CTCAGCAGGATTGATAAGACTACATTGTTC-  
3') was designed to create a construct (Vcat( G1172-G1191)) which truncated the C-  
terminus to residue D1171. Another oligo (5'-  
GAATTTGTCCCCTACAAGGAAGCTCCTGAAGATCTG-3') was designed to delete  
the central 50 residues (residues T940-E989) of the insert kinase domain, based on a  
sequence alignment with FGFR1 (Mohammadi et al, 1996). Sequence analysis  
detected an inadvertent Glu990-Val mutation. All DNA modification and restriction  
enzymes were purchased from New England Biolabs and oligonucleotides were  
purchased from Genosys Biotechnology.

The VEGFR2 50 construct was made in several steps to combine the necessary  
mutations into the baculovirus expression vector pAcSG2 (Pharmingen San Diego, CA). Step  
1; the coding region for Vcyt was PCR subcloned from the pET24a vector into the NcoI-KpnI  
sites of vector pAcSG2. Step2; a 2358bp Scal-BglII DNA fragment from plasmid pMGH4-Vcat  
( T940-E989,E990V) was ligated to a 1695bp BglII-ScaI DNA fragment from pMGH4-Vcat  
( G1172-G1191) creating a pMGH4-Vcat ( T940-E989,E990V, G1172-G1191) vector. Step 3; a  
913bp BstEII-EagI DNA fragment a pMGH4-Vcat ( T940-E989,E990V, G1172-G1191) was

ligated to a 3290bp *EagI*-*BstEII* DNA fragment from pAcSG2-Vcyt creating pAcSG2-Vcyt (T940-E989, E990V, G1172-G1191), also referred to as VEGFR2 50. This final construct was sequenced verified through the entire coding region and confirmed to contain only these known mutations from the wild-type sequence (sequence shown in Figure 1).

DNA encoding VEGFR2 50 was transfected into Sf9 cells with linearized baculovirus DNA according to the protocol of the manufacturer (Pharmingen San Diego, CA). Single plaques were isolated from this transfection and high titer stocks generated. All stocks were examined by isolation of baculoviral DNA and PCR amplification of the insert using the polyhedron forward and reverse primers (Invitrogen). Sf21 cells were infected at 1-1.5 million cells/mL at MOI=5 for 72 hours and harvested by centrifugation.

#### Purification Of VEGFR2 50 From Sf21 Cells

Cell pellets were lysed by dounce homogenization and sonication in 20 mM Tris pH 8.0, 20 mM NaCl, 5 mM DTT, and 5% (v/v) glycerol. The lysate was centrifuged for 50 minutes at 35,000 rpm in a Ti45 rotor. The soluble fraction was loaded onto a 40 ml Q-30 anion exchange column (Pharmacia) and eluted with a 20 mM to 600 mM NaCl gradient in 20 mM Tris pH 8.0, 5 mM DTT, and 5% (v/v) glycerol over 20 column volumes. VEGFR2 50 protein was pooled by SDS-PAGE gel analysis and by the presence of kinase activity as measured against gastrin substrate peptide substrate (Boehringer Mannheim). Pooled material was loaded onto a 40 mL hydroxyapatite (Bio-Rad) column and washed extensively with 20 mM Tris pH 8.0, 50 mM NaCl, 5 mM DTT, and 5% glycerol. Protein was eluted using a 500 mL linear gradient from 0 to 50 mM potassium phosphate pH 8.0, 50 mM NaCl, 5 mM DTT, and 5% glycerol. VEGFR2 50 protein was pooled by SDS-PAGE gel analysis and by the presence of kinase activity as measured against the gastrin peptide. Material from this column was then diluted 1:1 with 20 mM Tris pH 8.0, 20 mM NaCl, 5 mM DTT, and 5% glycerol and loaded onto an 8 mL Q-15 anion exchange column (Pharmacia). Protein was eluted using with a 180 mL linear NaCl gradient (20 mM-175 mM) in 20 mM Tris pH 8.0, 5 mM DTT, and 5% glycerol. VEGFR2 50 protein was pooled as described above. 4M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> was added to the pool to final concentration of 0.6

M and the pool loaded onto a 10 mL HP-phenyl sepharose column (Pharmacia). VEGFR2 50 protein was eluted using a 200 mL linear reverse gradient from 0.6 M to 0 M  $(\text{NH}_4)_2\text{SO}_4$  in 20 mM Tris and 5 mM DTT. Purified VEGFR2 50 protein was buffer exchanged into 50 mM Hepes pH 7.5, 10 mM DTT, 10 % glycerol, and 25 mM NaCl over a 500 ml G-25 column (Pharmacia) and concentrated to 1 mg protein/mL through a 10 kD cutoff polysulfone membrane (Amicon). Final material was aliquoted and flash frozen in liquid  $\text{N}_2$  and stored at  $-70^\circ\text{C}$ .

#### Kinetic Assays

The coupled spectrophotometric assays were done with purified VEGFR2 50 protein that was autophosphorylated under conditions: protein (4 mM), ATP (3 mM),  $\text{MgCl}_2$  (40 mM), DTT (5 mM), in Hepes (100 mM), 10% glycerol, pH 7.5 at  $4^\circ\text{C}$  for 1 hour.

#### Coupled Spectrophotometric Assay for the Forward Direction

Tyrosine kinase assays were monitored using a Beckman DU 650 Spectrophotometer. Production of ADP was coupled to oxidation of NADH using phosphoenolpyruvate (PEP) through the actions of pyruvate kinase (PK) and lactic dehydrogenase (LDH). The oxidation of NADH was monitored by following the decrease in absorbance at 340 nm ( $\epsilon_{340}=6.22 \text{ cm}^{-1} \text{ mM}^{-1}$ ). Typical reaction solutions contained: 1 mM PEP, 250 mM NADH, 50 units of LDH/mL, 20 units of PK/mL, 5 mM DTT, in 200 mM Hepes, pH 7.5 and varying concentrations of poly(E4Y1) (Sigma), ATP and  $\text{MgCl}_2$ . Assays were initiated with 40 nM of VEGFR2 50 protein.

#### Coupled Spectrophotometric Assay for the Reverse Reaction

ATP generation was coupled to production of NADH via the action of hexokinase (HK) and glucose-6-phosphate dehydrogenase (G6PD). In this assay, HK catalyzes the conversion of ATP to ADP and glucose-6-phosphate. Glucose-6-

phosphate is then oxidized to D-6-phosphogluconopyranose-1,5-lactone by G6PD with concomitant reduction of NAD to NADH which can be monitored at 340 nm. Typical assay solution contained: glucose (10 mM), NAD (40 mM), DTT (5 mM), MgCl<sub>2</sub> (4 mM), HK (15 unit/mL), G6PD (15 units/mL) and indicated concentrations of ADP and phospho-poly(E<sub>4</sub>Y). The reactions were initiated with addition of VEGFR2 50 protein (600-900 nM).

#### Evaluation of Potential Agonists and Antagonists of the VEGFR2 50 Protein

Based on the above spectrophotometric and kinetic assays, one can evaluate potential candidate agonists or antagonists of the VEGFR2 50 protein by addition of the candidate compounds to the above assay in a competition. As stated above, the kinetics of the activity of the VEGFR2 50 protein were measured against the gastrin peptide. The activity in the presence and absence of a candidate compound is measured and the resulting kinetic data is compared. The affinity of the candidate for the receptor will be reflected in the shift to the right of the kinetic curves indicating a competitive antagonist or with a decrease in the maximum activity, which would indicate a non-competitive antagonism. Conversely, a shift to the left of the kinetic curves would indicate a competitive agonist to the VEGFR2 50 protein. See generally, Bourne, H.R., et al. in, (1987) Basic & Clinical Pharmacology (Katzung, et al., eds) (Ch. 3) 9-22.

#### In Vitro Autophosphorylation Of VEGFR2 50 For Crystallization And Mass Spectrometry.

Aliquots of frozen VEGFR2 50 protein were thawed by immersion in cold H<sub>2</sub>O and pooled at 4°C. MgCl<sub>2</sub> and ATP were added to 26 mM and 4 mM, respectively. VEGFR2 50 was incubated at 4°C for 1 hour. This material (VEGFR2 50P) was then buffer exchanged into a solution of 10 mM Hepes 7.5, 10 mM DTT, and 10 mM NaCl and concentrated using a Centriprep-10 (Amicon) to 5 mg protein/mL.

#### Mass spectrometry

*Trypsin digestion.* Trypsin digestions of purified VEGFR2 50 and VEGFR2 50P were conducted at 37°C using 0.37 mg/mL protein in 25 mM  $\text{NH}_4\text{HCO}_3$  at pH 7.7 with a reaction volume of 100  $\mu\text{L}$  for two days.

*MALDI/MS.* MALDI-MS analyses were performed in a Voyager-Elite, time-of-flight mass spectrometer with delayed extraction (PerSeptive Biosystems, Inc., Framingham, MA). A volume of 1  $\mu\text{L}$  of digested protein sample was mixed with 1  $\mu\text{L}$  of matrix (a-cyano-4-hydroxy-cinnamic acid) in a solution of 50% (v/v) solution of acetonitrile and 0.25% (w/w) trifluoroacetic acid in water. Samples were irradiated with a nitrogen laser operated at 337 nm.

*NanoESI-MS.* NanoESI-MS analyses were performed on a triple quadrupole mass spectrometer (PE Sciex API III, Alberta, Canada) modified with a NanoESI source from Protana A/S, (Denmark). The ESI voltage was set at 700 V and the orifice settings were maintained at 100 V. 3  $\mu\text{L}$  of digested protein was mixed with 7  $\mu\text{L}$  of methanol and 0.5  $\mu\text{L}$  formic acid and then 4  $\mu\text{L}$  of this sample was injected into the mass spectrometer. Ion scans were used to obtain the sequence of phosphopeptides.

#### Crystallization and Data Collection

Purified phosphorylated VEGFR2 50 was concentrated on average to 5 mg protein/mL using a Centricon-10 centrifugal concentrator. Crystals were grown by the hanging drop vapor diffusion method at 4°C. Drops containing 2  $\mu\text{L}$  of protein solution and 2  $\mu\text{L}$  of a mother liquor solution (100 mM Hepes at pH 7.2, 2 M  $(\text{NH}_4)_2\text{SO}_4$ , and 2% (v/v) monomethylether polyethylene glycol mW=550) were equilibrated above a 1 mL reservoir of the mother liquor solution to which 50 mM  $\beta$ -mercaptoethanol had been added. Crystals appeared after 3-4 days and grew to as large as 0.3 x 0.2 x 0.5 mm over 21 days.

X-ray diffraction data sets were collected using a Rigaku RU-200 rotating anode X-ray generator (CuK $\alpha$ ) operated at 50 kV and 100 mA and equipped with Supper focusing mirrors and a MAR345 MAR Research image plate detector. Data

5 collection on frozen crystals was done by transferring a crystal into a cryoprotectant  
solution (100 mM Hepes at pH 7.2, 2.2 M  $(\text{NH}_4)_2\text{SO}_4$ , 0.6 M sucrose, 0.55 M  
glucose, and 2% (v/v) monomethylether polyethylene glycol MW=550), flash freezing  
10 the crystal in liquid nitrogen, and then transferring the frozen crystal into a stream of  
5 nitrogen at -186 °C. Data was integrated and scaled using DENZO and SCALEPACK  
(Otwinowski, 1993) Data collection statistics are given in Table 2.

15 Initial protein phases were obtained using the AMoRe molecular replacement  
program (Navaza, 1994), molecule 1 of the FGFR1 structure (Mohammadi et al.,  
20 1996; PDB entry 1FGK) as a search probe, and the native1 data set. The correct  
solution was achieved by including the FGFR1 sidechains and removing from mobile  
20 residues of the activation loop (640-660), the N-terminus (464-467), a short loop  
(517-520), and the C-terminus (760-762) from the search model. The correct solution  
15 was the top peak in the rotation and translation functions with a correlation coefficient  
of 0.31. Rigid body refinement in AMoRe improved the solution to a correlation  
25 coefficient of 0.49 and an *R*-factor of 46.3% in the 12.0 - 4.0 Å resolution range. The  
correctness of this solution was cross-checked by calculation of a difference Fourier  
with a  $\text{KAu}(\text{CN})_2$  derivative. This derivative was generated by soaking a crystal for 3  
30 days in reservoir solution containing 0.5 mM  $\text{KAu}(\text{CN})_2$  and then increasing the heavy  
20 atom concentration to 5 mM and soaking for an additional 64 hours. Scaling of data  
sets, Patterson calculations, Fourier calculations, and the generation of phases were  
done using Xtalview (McRee et al., 1992)

35 Refinement of the model was done using Xplor version 3.1 (Brünger, 1992).  
25 Calculation of electron density maps and model fitting was done using XtalView  
(McRee et al., 1992) Refinement was begun using a data set collected at 4 °C  
40 (native2) and was completed using a data set (native3) collected at -186 °C. The  
final *R*-factor is 20.2% for data in the range 8-2.4 Å ( $F_o > 2\sigma$ ). The average *B* value  
for all atoms is 31.8 Å<sup>2</sup> for protein atoms and 42.8 Å<sup>2</sup> for water molecules. The final  
45 30 model includes residues 820-939, 998-1047, and 1064-1168; of these residues the  
sidechains of K838, R842, F845, K939, D998, K1023, R1027, Y1038, K1039, K1110,  
and E1113 could not be modeled beyond C due to a lack of interpretable density.  
Analysis of main-chain torsion angles as done using PROCHECK (Laskowski et al.,

1993) shows of the 275 residues in the model none occur in the disallowed region and only 4 occur in the generously allowed region of a Ramachandran plot. 182 water molecules were fit to electron density peaks which were greater than 3 $\sigma$  and were located in positions to make reasonable hydrogen bonds to the protein or other water molecules.

Superpositions of various kinase structures was done using the graphics program Insight II (Molecular Simulations Inc, San Diego, CA).

#### Example 1- Structure Determinations

The tyrosine kinase domain of human VEGFR-2 lacking the 50 central residues of the 68 residues of the KID was expressed in a baculovirus/insect cell system. Of the 1356 residues of full-length VEGFR-2 this construct (VEGFR2 50) contains residues 806-939 and 990-1171 of the cytosolic domain (Figure 1). VEGFR2 50 also contains one point mutation (E990V) within the KID relative to wild-type VEGFR-2.

In addition to catalyzing its autophosphorylation, VEGFR2 50 is also capable of catalyzing phosphorylation of a poly(E4Y) exogenous substrate. Detailed kinetic analysis (Table 1) revealed that its kinetic parameters were nearly identical to that of a comparable VEGFR-2 protein construct containing the entire KID (Parast et al., in press). These results taken together indicate that VEGFR2 50 is a fully active functional enzyme. Therefore, deletion of 50 central residues of the KID has no observed effect on the catalytic steps of the phosphotransfer reaction. It was also determined that deletion of more than 60 amino acids from the KID region did cause a diminishment in the activity of the enzyme.

Table 1: Kinetic constants of VEGFR2 50

Forward Reaction			
Substrate	$K_M$ (mM)	$k_{cat}$ ( $s^{-1}$ )	$k_{cat}/K_M$ ( $s^{-1}M^{-1}$ )
MgATP	0.153	13.3	$87 \times 10^3$
poly(E4Y)	2.1		$63 \times 10^2$
Mg <sup>2+</sup>	6.8		$20 \times 10^2$
Reverse Reaction			
Substrate	$K_M$ (mM)	$k_{cat}$ ( $s^{-1}$ )	$k_{cat}/K_M$ ( $s^{-1}M^{-1}$ )
MgADP	0.056	0.13	$23 \times 10^2$
P-poly(E4Y)	1.0		$13 \times 10^1$

The VEGFR-2 KID sequence is hydrophilic and highly charged, containing 6 lysine, 5 arginine, 8 glutamic acid, and 5 aspartic acid residues (Figure 1). Initially several protein constructs containing the VEGFR-2 catalytic domain with the entire KID were generated. After exhaustive attempts to crystallize these protein constructs failed to yield even marginal crystals, the VEGFR2 50 construct was created to test the idea that the highly charged KID was interfering with crystallization. As determined by dynamic light scattering this VEGFR2 50 construct, which eliminated 14 charged residues, exhibited markedly better stability to temperature and protein concentration than protein constructs containing the entire KID.

For crystallization, purified VEGFR2 50 was autophosphorylated in vitro by incubation with MgATP. Matrix-assisted laser desorption ionization (MALDI) and

nanoelectrospray ionization (NanoESI) mass spectrometry analysis of full-length phosphorylated VEGFR2 50 (VEGFR2 50P) protein and tryptically digested peptides indicates phosphorylation of Y1059 using the autophosphorylation conditions described here. Crystals diffracting to 2.2 Å were obtained of VEGFR2 50P in an unligated state. The crystals belong to the orthorhombic space group P2<sub>1</sub>2<sub>1</sub>2<sub>1</sub> with one VEGFR2 50P molecule in the asymmetric unit. Initial crystallographic phases were determined by molecular replacement using the structure of the unphosphorylated kinase domain of FGFR1 (Mohammadi et al., 1996) as a search model. The correctness of the molecular replacement solution was cross-checked using a gold cyanide derivative. The derivative data, however, was not used for phase calculations of electron density maps used to build the model. The structure has been refined to an *R*-factor of 20.2% for 8-2.4 Å data (*F*<sub>o</sub> > 2σ). VEGFR2 50P residues for which backbone atoms were not modeled due to disorder include the N-terminal residues 806-819, residues 990-997 of the KID, residues 1048-1063 of the activation loop, and residues 1169-1171 of the C-terminus. Structure determination statistics are included in Table 2.

**Table 2: VEGFR2 50P structure determination statistics**

Data Set	Native (3)	Native (1)	Native (2)	KAu(CN) <sub>2</sub>
Data resolution (Å)	15-2.2	20-3.0	15-2.4	15.3.1
<i>R</i> <sub>sym</sub> (%)	5.2 <sup>a</sup> {19.6} <sup>b</sup>	8.4 (19.2)	7.0 (21.9)	7.1 (19.5)
Completeness (%)	93.0 (81.0)	97.5 (98.4)	98.8 (98.8)	96.5 (95.0)
Temperature (°C)	-186	room (~21)	4	4
Unit cell a (Å)	95.41	97.10	98.52	97.71
Unit cell b (Å)	96.04	96.94	96.50	96.97
Unit cell c (Å)	38.22	38.63	38.56	38.52
Refinement resolution (Å)	8-2.4	--	--	--
Refined <i>R</i> (%)	20.2 <sup>c,d</sup>	--	--	--

<sup>a</sup>  $R_{\text{sym}} = \frac{\sum hkl \sum i |I_i(hkl) - \langle I(hkl) \rangle|}{\sum hkl \sum i I_i(hkl)}$

<sup>b</sup> Value in parenthesis is for highest (resolution shell)

$$R = \frac{\sum_{hkl} |F_O(hkl)| - |F_C(hkl)|}{\sum_{hkl} |F_O(hkl)|}$$

where  $F_O$  and  $F_C$  are the observed and calculated structure factors, respectively ( $F_O > 2$ )

<sup>d</sup> Model includes 275 protein residues and 182 water molecules

### Overall Kinase Fold

Analogous to previously reported structures of both serine/threonine and tyrosine protein kinases, VEGFR2 50P is folded into two lobes with catalysis of phosphotransfer taking place in a cleft between the two lobes (reviewed in Cox et al., 1994; Johnson et al., 1996). A C $\alpha$  trace of the VEGFR2 50P structure is shown in Figure 2a. Kinase secondary structural elements are designated (Figure 1) according to the convention originally given for cAPK (Knighton et al., 1991). The N-terminal lobe (approximately residues 820-920) folds into a twisted  $\beta$ -sheet with one  $\alpha$ -helix (C). The structure comprises five antiparallel strands (1-5), three of which (1-3) are highly curved and curl over the other two strands (4-5). The larger C-terminal domain (approximately residues 921-1033) contains two antiparallel  $\beta$ -strands (7-8), which lie at the top of the C-terminal domain adjacent to the N-terminal  $\beta$ -sheet. Seven  $\alpha$ -helices (D, E, E-F, G, H, I) form the remaining core of the C-terminal domain. Like other kinases, VEGFR2 50P contains two functionally important loop regions: the glycine-rich nucleotide binding loop (residues 841-846), the catalytic loop (residues 1026-1033) and the activation loop (residues 1046-1075) (Figures 1 and 2a).

Of the reported kinase structures, the VEGFR2 50P structure resembles most closely that of the catalytic domain of FGFR1 (Mohammadi et al., 1996; PDB entry 1FGK) with which it shares approximately 55% sequence identity (Figure 1). Since the two molecules in the crystallographic asymmetric unit of the FGFR1 structure solution are very similar, comparisons to VEGFR2 50P will primarily be described only for FGFR1 molecule A. Least squares superposition of 82 C $\alpha$  positions of (1-5) of the N-terminal lobe or 152 C $\alpha$  positions residues (D, E, F, G, H, I) of the C-terminal lobe between FGFR1 and VEGFR2 50P result in respective rms deviations of 0.40 Å and

0.52 Å. A relative rotation of approximately 5° between the two lobes results in the interlobe cleft of VEGFR2 50P being slightly larger and more open. Measurement of distances between equivalent C's (K523 and R675 of FGFR1, S877 and R1080 of VEGFR2 50P) at the ends of the cleft reveal that this distance is 25.3 Å in VEGFR2 50P as compared to 23.2 Å in FGFR1. This is however a minor difference, as compared to much larger relative lobe rotations observed among kinase structures in various ligation and phosphorylation states (Johnson et al., (1996) *Cell* 85, 149-158). For example, the inter-lobe orientation seen here for VEGFR2 50P is in an approximately 20° more open conformation than that seen in the ternary complex structure of the phosphorylated kinase domain of IRK bound to the ATP analog AMP-PNP and a peptide substrate (Hubbard, (1997) *EMBO J.* 16, 5572-5581; PDB entry 1IR3) (Figure2c).

While the  $\alpha$ -strand positions of the N-terminal lobe agree well between VEGFR2 50P and FGFR1, the structures do diverge significantly at the N-terminal residues preceding the first conserved region starting at residue W827 (Figure 2a and 2b). The first 14 residues (M806-E819) of VEGFR2 50P are completely disordered and the next seven residues (L820-R826) form an extended loop structure. It is likely that residues 806-819 do not form part of the active kinase region but are instead part of, or are adjacent to, the juxtamembrane region of VEGFR-2. Residues 820-826 do seem to be part of the kinase domain, although a flexible one, as analogous residues are also ordered in the structures of FGFR1, IRK, and the non-receptor tyrosine kinase Lck (Yamaguchi and Hendrickson, (1996) *Nature* 384, 484-489). Other differences between the VEGFR2 50P structure and other kinase structures occur at the kinase insert domain and the activation loop (discussed below).

#### Catalytic loop and ATP binding site

In protein kinases, the loop between E and 7 has been termed the catalytic loop as it contains an invariant aspartic acid (D1028) that is believed to function as a catalytic base in the phosphotransfer reaction (Johnson et al., 1996). This aspartic acid is part of a stretch of residues (H1026-N1033) whose sequence HRDLAARN is highly conserved among protein tyrosine kinases. In VEGFR2 50P the backbone

5 position and most sidechain positions of this loop are similar to those in the  
unliganded FGFR1 and ternary phosphorylated IRK (IRKP) complex structures. As  
seen in these previous structures the sidechain carboxylate of the catalytic loop  
10 aspartic acid (D1028) is hydrogen bonded to the sidechains of the conserved arginine  
5 (R1032) and asparagine (N1033) (Figure 3).

The ATP binding site of protein kinases lies at the cleft between the N and C-  
15 terminal lobes (Figure 2c). For VEGFR2 50P, the residues forming this site consist  
primarily of residues E917-N923, joining the two lobes, and residues L840-I849 which  
10 include parts of 1, 2, and the glycine-rich loop of G841-G846. The glycine-rich loop,  
also referred to as the nucleotide binding loop, is a flexible segment whose position  
20 differs among kinase structures in various activated and liganded states. In  
VEGFR2 50P this loop is fairly well ordered and all atoms could be modeled with the  
exception of the sidechains of R842 and F845. The relative position and  
15 conformation of this loop is similar to that observed in the unliganded FGFR1 structure.  
However, this position is markedly different from that in the IRKP ternary complex  
25 structure in which the approximately 20° relative rotation of the N and C-terminal  
lobes results in the glycine-rich loop being 5 Å closer to the C-terminal lobe than in  
VEGFR2 50P structure.

20 In reported kinase structures with bound ATP or an ATP analog, the adenine  
ring makes two conserved hydrogen bonds with the protein backbone. In the  
35 structure of FGFR1 with AMP-PCP bound (Mohammadi et al., 1996) these hydrogen  
bonds are between the adenine NH<sub>2</sub> and the backbone C=O of E562 (E917  
25 VEGFR2 50P) and between the adenine N1 and the backbone NH of A546 (C919  
VEGFR2 50P). Although the structure presented here does not contain a bound  
40 nucleotide, the similarities in the positions of these backbone atoms to those in  
FGFR1 indicate that these hydrogen bonds would be formed in a VEGFR2 50P-ATP  
complex and therefore the adenine is expected to bind in a similar position (Figure 4).

30 Variation in the ATP-binding sites of kinases involved in disease is of  
considerable importance in the design of selective ATP-competitive inhibitors as  
45 therapeutics. A comparison of the ATP binding sites of FGFR1 and VEGFR2 50P

reveals that while the overall architecture of the site is conserved, several sequence differences result in differences in the shape of the accessible area for ligand binding. Specific sequence differences between FGFR1 and VEGFR-2 in this site include: V899 (I545 FGFR1), F918 (Y563 FGFR1), C919 (A564 FGFR1), and C1045 (A640 FGFR1) (Figure 4). Similarly, comparison to the ternary IRKP complex structure reveals variation in the adenine site at V916 (M1076 IRK), F918(L1078), C919 (M1079 IRK), L1035 (M1139 IRK), and C1045 (G1149 IRK). Even greater sequence and structural variation in the adenine site is seen when the VEGFR2 50P structure is compared to serine/threonine kinase structures, suggesting that these differences are useful in the design of selective ATP-competitive inhibitors.

#### Activation loop

Protein kinases contain a large flexible loop, referred to as the activation loop (A-loop) whose conformation is postulated to regulate kinase activity (Figure 2). In many kinases the conformation of the AL is controlled by the phosphorylation of specific A-loop residues (Johnson et al., 1996). The loop can be generally defined as beginning with the conserved residues DFG and ending at the conserved APE sequence (Johnson et al., 1996). In VEGFR-2 this segment corresponds to D1046-E1075 and contains two tyrosines (Y1054 and Y1059). Both Y1054 and Y1059 were found to be autophosphorylation sites when the cytosolic domain of VEGFR-2 was expressed in *E. coli* (Dougher-Vermazen et al., 1994). Using the in vitro autophosphorylation protocol described here for VEGFR2 50, a stable phosphorylation site is indicated at Y1059, however no evidence of phosphorylation of Y1054 was detected.

In this unliganded VEGFR2 50P structure presented here, the A-loop appears quite mobile and interpretable electron density was not present for most of the central portion of the loop (G1048-G1063). This disorder is consistent with mobility of the A-loop deduced from other kinase structures. For example, of the two molecules in the asymmetric unit of the unphosphorylated FGFR1 kinase structure the center of the A-loop has relatively high temperature factors in molecule A and is completely disordered in molecule B. Although residues 1048-1063 could not be modeled in

5 VEGFR2 50P, unambiguous electron density was present for residues D1064-E1075,  
clearly indicating that these residues adopt a conformation similar to that observed in  
the unphosphorylated FGFR1 structure. The segment of D1064-P1068 has an  
10 extended structure that lies adjacent to the catalytic residues D1028 and R1032  
(Figure 3a). Comparison to the structure of the (MgAMP-PNP)-peptide-IRKP  
complex structure indicates that the position of R1066-P1068 in this VEGFR2 50P  
structure is inhibitory to substrate binding. P1066 occupies equivalent space  
15 allocated to the tyrosine sidechain of the peptide substrate in the ternary IRK3P  
complex structure. The conformation of residues L1069-E1075 is similar to that in the  
10 ternary IRKP complex structure, however there is a complete directional change at  
P1068 (P1172 IRK) between the two structures. In the IRK structure residues N-  
20 terminal to this proline are directed toward EF while in VEGFR2 50P they are directed  
toward D on the opposite side of the protein (Figures 2 and 3).

15 Despite the phosphorylation of Y1059 prior to crystallization, the conformation  
seen here for residues D1064-P1068 is similar to the inhibitory conformation  
observed for analogous residues in the unphosphorylated FGFR1 structure. Y1059  
in VEGFR2 50 corresponds to a relatively conserved phosphorylation site among  
30 protein tyrosine kinases. In the ternary IRKP complex structure and the  
20 phosphorylated lymphocyte kinase (Lck) structure (Yamaguchi and Hendrickson,  
1996) the tyrosine at this position (Y1163 IRK, Y394 Lck) is phosphorylated and the  
A-loop has a non-inhibitory conformation similar to that observed in a phosphorylated  
35 cAPK ternary complex structure (Zheng et al., 1993). The interactions the phosphate  
group at this position makes with other protein residues are believed to help stabilize  
25 an A-loop conformation that allows substrate and ATP binding (Johnson et al., 1996;  
Hubbard, 1997). However, since this VEGFR2 50P structure described here does not  
40 exhibit a similar open A-loop conformation but rather has an inhibitory conformation  
with much of the loop disordered it is possible that the monophosphorylated A-loop of  
VEGFR2 50P exists in a dynamic equilibrium involving several conformations and that  
30 the conformation observed here is the one most favored in this crystal environment.

5

**Kinase Insert Domain:**

10

15

20

25

The kinase insert domain occurs in the kinase C-terminal lobe and connects helices D and E. In VEGFR-2 this region corresponds to a 68 residue stretch from N933 to L1000 (Figure 1). The lack of effect on intrinsic kinase activity (noted above) of deletion of residues T940-E989 is perhaps not surprising as the ends of the KID domain occur relatively far away (approximately 35-40 Å) from the catalytic site and on the opposite side of the protein from the position of the activation loop (Figure 2). These results are consistent with those for the CSF-1 receptor kinase in which deletion of 58 of the 64 residues of the CSF-1 KID only decreased its ability to phosphorylate a peptide substrate by 10% (Taylor et al., 1989). Deletion of the entire 98 residues of PDGFR, however, resulted in an 80% decrease in kinase activity towards a peptide substrate (Severinsson et al., (1990) *Mol. Cell. Biol.* 10, 801-809). Thus, the present invention allows for the production of a synthetic catalytic linker which recognizes that the majority of KID is not required for catalysis but rather only a small number of residues must be present to form a linker between  $\alpha$ D and  $\alpha$ E so as to maintain a competent kinase structure.

30

35

40

In the VEGFR2 50P structure following D, residues N933-P937 form a loose turn and an extended strand whose ends are roughly perpendicular to the axes of D and I at the C-terminus. In different Fourier maps, the electron density is strong and clear for residues N933-P937 and becomes weak for Y938 and K939 (sidechains of Y938 and K939 are not modeled) (Figure 5). The 50 residue deletion in VEGFR2 50 directly follows K939 so that the residue immediately C-terminal to K939 is V990, maintaining the residue numbering in full-length VEGFR-2. Residues V990-K997 are disordered and interpretable electron density begins again at D998. Residues D998-T1001 then form a short strand that joins E at residue L1002 (Figures 5 and 6).

45

50

The two strands at the N-terminal and C-terminal ends of the KID form a pseudo two-stranded parallel  $\beta$ -sheet structure that is different from the conformations seen in this region of other kinase structures. The two ends of the KID thus make a variety of interactions which may help to stabilize the overall conformation and position of this domain in VEGFR-2. The sidechain of K931 makes an ionic

55

5 interaction with the sidechain of E934 and also makes a hydrogen bond to the  
backbone carbonyl of D998 (Figure 6). Hydrogen bonding interactions between the  
strands include: E934 backbone C=O to L1000 NH, V936 NH to L1000 C=O, and  
10 P937 C=O to L1002 NH. In addition to these polar interactions, the sidechains of  
5 F935, P937, and L1000 are involved in extensive hydrophobic contacts. The  
sidechain of F935 is nestled in a hydrophobic pocket formed by the sidechains of  
L928, P937, L1000, L1002, L1005, L1101, and Y1130 (Figures 5 and 6). The L1000  
15 sidechain also packs against the sidechains of Y927, K931, H1004, and Y1008.

10 It has been found by the applicants that deletion of portions of the KID also  
impart other useful and desirable characteristics to the modified VEGFR-2  
20 polypeptide. The modified polypeptide has exhibited greater stability when exposed to  
higher temperatures in solution than the wild-type protein. Additionally, the modified  
polypeptide has also exhibited improved solubility than the wild-type protein. It is  
25 apparent to those skilled in the art that these properties allow improvements in  
various commercial aspects of the present invention. Examples of potential uses for  
the modified proteins include high-throughput screening of potential ligands for the  
receptor by various methods including those based on gene-chip technology  
30 (Affymax, Inc.) phage-display peptide libraries (The Ph.D. Kit® by New England  
20 BioLabs, Inc.) as well as in-depth analysis via FT-NMR.

35 It is therefore contemplated that the entire KID can be deleted and retain  
some catalytic activity in other related RTKs such as but not limited to PDGF and  
and other previously mentioned proteins. Furthermore, in one embodiment of the  
25 invention the entire KID is deleted and replaced with a synthetic catalytic linker of at  
least one amino acids such that both the catalytic activity and the crystallizability of  
40 the protein is retained.

#### Cloning of The PDGFR Protein

45 30 In this example, the PDGFR polypeptide is cloned using the methods outlined  
for VEGFR-2 above. The coding sequence for PDGFR is derived from the sequence  
disclosed by Matsui, T., et al., (1989) Science 243: 800-804 (Accession No. 66814).

5

PCR oligonucleotide primers are then made which code for residues located in the cytoplasmic domain and the catalytic domain of the protein. The catalytic domain of PDGFR is shown starting at residue 689 (N) and ending at residue 791 (T) in Figure 1.

10

5

The remainder of the cloning and purification steps would be similar to those disclosed for the VEGFR2 50 protein and use technology well known to those skilled in the art.

15

10

It is contemplated that other members of the RTK family and other uses for the data disclosed herein and are not limited by the examples shown.

20

15

25

30

35

40

45

50

55

## Claims

5

10

15

20

25

30

35

40

45

50

55

5

## WHAT IS CLAIMED IS:

10

1. A method for assaying a candidate compound for its ability to interact with a modified receptor tyrosine kinase (RTK) polypeptide comprising:

15

- a) expressing an isolated DNA sequence or variants thereof encoding the modified RTK gene construct wherein said RTK gene contains a synthetic catalytic linker wherein said linker comprises at least one amino acid from the kinase insert domain (KID) of the VEGFR-2 gene catalytic region, in a host capable of producing a form of the polypeptide which form may be assayed for interaction of said polypeptide with said candidate substance;

20

- b) exposing said modified polypeptide to said candidate substance; and

25

- c) evaluating the interaction of said polypeptide with said candidate substance.

2. The method of claim 1, wherein said evaluation step further comprises:

30

- (a) crystallizing said modified polypeptide in a condition suitable for x-ray crystallography; and
- (b) conducting said x-ray crystallography on said polypeptide.

35

3. A method for assaying a candidate compound for its ability to interact with a modified VEGFR-2 receptor polypeptide comprising:

40

- a) expressing an isolated DNA sequence or variants thereof encoding the modified VEGFR-2 gene construct wherein said VEGFR-2 gene contains a synthetic catalytic linker wherein said linker comprises at least one amino acid from the kinase insert domain (KID) of the VEGFR-2 gene catalytic region, in a host capable of producing a form of the polypeptide which form may be assayed for interaction of said polypeptide with said candidate substance;

45

- b) exposing said modified polypeptide to said candidate substance; and

50

55

5

c) evaluating the interaction of said modified polypeptide with said candidate substance.

10

4. The method of claim 2, wherein said evaluation step further comprises:  
(a) crystallizing said modified polypeptide in a condition suitable for x-ray crystallography; and

15

(b) conducting said x-ray crystallography on said polypeptide.

20

5. An isolated DNA sequence or variants thereof encoding a modified RTK gene construct wherein said RTK gene contains a synthetic catalytic linker wherein said linker comprises at least one amino acid from the kinase insert domain of the RTK gene catalytic region.

25

6. An isolated DNA sequence or variants thereof encoding a modified VEGFR-2 gene construct wherein said VEGFR-2 gene contains a synthetic catalytic linker wherein said linker comprises at least one amino acid from the kinase insert domain of the VEGFR-2 gene catalytic region.

30

7. The isolated oligonucleotide sequence of claim 6 comprising a DNA sequence or variants thereof in SEQ. ID NO. 5.

35

8. The isolated oligonucleotide sequence of claim 6 comprising a DNA sequence or variants thereof in SEQ. ID NO. 6.

40

9. A method of assessing compounds which are agonists or antagonists of the activity of the a modified RTK gene polypeptide wherein said modified RTK gene contains a synthetic catalytic linker wherein said linker contains at least one amino acid from the kinase insert domain of the RTK polypeptide catalytic region comprising:

45

a) crystallizing said modified RTK polypeptide;

50

55

5

b) obtaining crystallography coordinates for said crystallized modified RTK polypeptide;

10

c) applying said crystallography coordinates for said modified RTK polypeptide to a computer algorithm such that said algorithm will generate a model of said RTK polypeptide suitable for use in designing molecules that will act as agonists or antagonists to said polypeptide; and

15

d) applying an iterative process whereby various molecular structures are applied to said computer generated model to identify potential agonists or antagonists to said polypeptide.

20

10. A method of assessing compounds which are agonists or antagonists of the activity of the a modified VEGFR-2 gene polypeptide wherein said modified VEGFR-2 gene contains a synthetic catalytic linker wherein said linker comprises at least one amino acid from the kinase insert domain of the VEGFR-2 polypeptide catalytic region comprising:

25

30

a) crystallizing said modified VEGFR-2 polypeptide;

b) obtaining crystallography coordinates for said crystallized modified VEGFR-2 polypeptide;

35

c) applying said crystallography coordinates for said modified VEGFR-2 polypeptide to a computer algorithm such that said algorithm will generate a model of said VEGFR-2 polypeptide suitable for use in designing molecules that will act as agonists or antagonists to said polypeptide; and

40

d) applying an iterative process whereby various molecular structures are applied to said computer generated model to identify potential agonists or antagonists to said polypeptide.

45

50

55

- 5                    11. The method of claim 10, wherein said modified VEGFR-2 polypeptide  
comprises the VEGFR2 50 polypeptide of Seq. ID No. 5.
- 10                   12. An isolated DNA sequence comprising a DNA sequence or variants thereof  
encoding the VEGFR-2 gene construct having the x-ray coordinates of Figure  
5.
- 15                   13. A method for preparing proteins or polypeptides of the receptor tyrosine  
kinase family such that they are suitable for measurement by x-ray  
crystallography comprising:
- 20                   a) identification of the Kinase Insert Domain within the catalytic domain of said  
proteins;
- 25                   b) deletion of a specific number of amino acid residues from said Kinase  
Insert Domain such that the modified polypeptide now has a stable  
conformation such that it may form a crystalline state suitable for being  
measured by x-ray crystallography; and
- 30                   c) crystallizing said modified polypeptide.
- 35                   14. A process of drug design for compounds which interact with RTK polypeptides  
comprising:
- 40                   a) deletion of a portion of the KID of the target RTK polypeptide;
- 45                   b) crystallizing said target RTK polypeptide;
- 50                   c) resolving the x-ray crystallography of said target RTK polypeptide;
- 55                   d) applying the data generated from resolving the x-ray crystallography of said  
target RTK polypeptide to a computer algorithm which will generate a model of  
said target RTK polypeptide suitable for use in designing molecules that will

- 5 act as agonists or antagonists to said polypeptide; and
- e) applying an iterative process whereby various molecular structures are  
10 applied to said computer generated model to identify potential agonists or  
antagonists to said target RTK polypeptide.
- 15 15. A process of drug design for compounds which interact with modified VEGFR-  
2 polypeptides comprising:
- a) deletion of a portion of the KID of the modified VEGFR-2 polypeptide;
- 20 b) crystallizing said modified VEGFR-2 polypeptide;
- c) resolving the x-ray crystallography of said modified VEGFR-2 polypeptide;
- 25 d) applying the data generated from resolving the x-ray crystallography of said  
modified VEGFR-2 polypeptide to a computer algorithm which will generate a  
model of said modified VEGFR-2 polypeptide suitable for use in designing  
30 molecules that will act as agonists or antagonists to said polypeptide; and
- e) applying an iterative process whereby various molecular structures are  
35 applied to said computer generated model to identify potential agonists or  
antagonists to said modified VEGFR-2 polypeptide.
- 40 16. The method of claim 15, wherein said modified VEGFR-2 polypeptide  
comprises the VEGFR2 50 polypeptide of Seq. ID No. 5.






Figure 2a



Figure 2b



Figure 2c




Fig. 58




Fig. 4




Fig. 5





Fig. 6



## SEQUENCE LISTING

<110> McTigue Ph.D., Michele A.

<120> Modifications of The VEGF Receptor-2 Protein And  
Methods Of Use

<130> Modified VEGFR-2 protein

<140> 0125-0016

<141> 1998-09-08

<160> 1

<170> PatentIn Ver. 2.0

<210> 1

<211> 367

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Human Aorta cDNA  
library

<400> 1

Met Asp Pro Asp Glu Leu Pro Leu Asp Glu His Cys Glu Arg Leu Pro  
1 5 10 15

Tyr Asp Ala Ser Lys Trp Glu Phe Pro Arg Asp Arg Leu Lys Leu Gly  
20 25 30

Lys Pro Leu Gly Arg Gly Ala Phe Gly Gln Val Ile Glu Ala Asp Ala  
35 40 45

Phe Gly Ile Asp Lys Thr Ala Thr Cys Arg Thr Val Ala Val Lys Met  
50 55 60

Leu Lys Glu Gly Ala Thr His Ser Glu His Arg Ala Leu Met Ser Glu  
65 70 75 80

Leu Lys Ile Leu Ile His Ile Gly His His Leu Asn Val Val Asn Leu  
85 90 95


Leu Gly Ala Cys Thr Lys Pro Gly Gly Pro Leu Met Val Ile Val Glu  
100 105 110

Phe Cys Lys Phe Gly Asn Leu Ser Thr Tyr Leu Arg Ser Lys Arg Asn

115	120	125
Glu Phe Val Pro Tyr Lys Thr Lys Gly Ala Arg Phe Arg Gln Gly Lys		
130	135	140
Asp Tyr Val Gly Ala Ile Pro Val Asp Leu Lys Arg Arg Leu Asp Ser		
145	150	155
Ile Thr Ser Ser Gln Ser Ser Ala Ser Ser Gly Phe Val Glu Glu Lys		
165	170	175
Ser Leu Ser Asp Val Glu Glu Glu Glu Ala Pro Glu Asp Leu Tyr Lys		
180	185	190
Asp Phe Leu Thr Leu Glu His Leu Leu Ile Cys Tyr Ser Phe Gln Val		
195	200	205
Ala Lys Gly Met Glu Phe Leu Ala Ser Arg Lys Cys Ile His Arg Asp		
210	215	220
Leu Ala Ala Arg Asn Ile Leu Leu Ser Glu Lys Asn Val Val Lys Ile		
225	230	235
Cys Asp Phe Gly Leu Ala Arg Asp Ile Tyr Lys Asp Pro Asp Tyr Val		
245	250	255
Arg Lys Gly Asp Ala Arg Leu Pro Leu Lys Trp Met Ala Pro Glu Thr		
260	265	270
Ile Phe Asp Arg Val Tyr Thr Ile Gln Ser Asp Val Trp Ser Phe Gly		
275	280	285
Val Leu Leu Trp Glu Ile Phe Ser Leu Gly Ala Ser Pro Tyr Pro Gly		
290	295	300
Val Lys Ile Asp Glu Glu Phe Cys Arg Arg Leu Lys Glu Gly Thr Arg		
305	310	315
Met Arg Ala Pro Asp Tyr Thr Thr Pro Glu Met Tyr Gln Thr Met Leu		
325	330	335
Asp Cys Trp His Gly Glu Pro Ser Gln Arg Pro Thr Phe Ser Glu Leu		
340	345	350
Val Glu His Leu Gly Asn Leu Leu Gln Ala Asn Ala Gln Gln Asp		
355	360	365

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/20319

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : Please See Extra Sheet. US CL : 117/70; 435/7.2, 69.1, 183, 194; 436/4, 86; 530/350 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 117/70; 435/7.2, 69.1, 183, 194; 436/4, 86; 530/350 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WEST, MEDLINE, CAPLUS		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	AL-OBEIDI et al. Protein Tyrosine Kinases: Structure, Substrate Specificity, and Drug Discovery. Biopolymers (Peptide Science). 1998, Vol. 47, pages 197-223, especially page 213, first column, second paragraph, to column 2, first paragraph.	1-4, 9-10
Y	WEI et al. Expression, Characterization, and Crystallization of the Catalytic Core of the Human Insulin Receptor Protein-tyrosine Kinase Domain. J. Biol. Chem. 07 April 1995, Vol. 270, No. 14, pages 8122-8130, especially page 8130, Figure 9.	2, 4, 9-11, 13-16
Y	SCOPES. Protein Purification: Principles and Practice. Second Edition, 1987, pages 297-301, see entire document.	2, 4, 9-11, 13-16
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:		*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance		*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*C* earlier document published on or after the international filing date		*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* documents which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)		*G* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search	Date of mailing of the international search report	
23 NOVEMBER 1999	27 JAN 2000	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer JOSEPH MURPHY 	
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196	

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/20319

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	MOHAMMADI et al. Structures of the Tyrosine Kinase Domain of Fibroblast Growth Factor Receptor in Complex with Inhibitors. Science. 09 May 1997, Vol. 276, pages 955-960, especially page 957, Figure 3.	1-2, 4, 9, 13-14 ----- 3, 10-11, 15-16
A	MCDONALD et al. The first structure of a receptor tyrosine kinase domain: a further step in understanding the molecular basis of insulin action. Structure. 15 January 1995, Vol. 3, No. 1 pages 1-6, especially page 4, Figure 4.	2, 4, 9-10, 13-16
A	HUBBARD et al. Crystal structure of the tyrosine kinase domain of the human insulin receptor. Nature. 29 December 1994, Vol. 372, pages 746-754, especially page 747, Table 1.	2, 4, 9-10, 13-16
X --- Y	SINGH et al. Structure-Based Design of a Potent, Selective, and Irreversible Inhibitor of the Catalytic Domain of the erbB Receptor Subfamily of Protein Tyrosine Kinases. J. Med. Chem. 1997, Vol. 40, No. 7, pages 1130-1135, especially page 1133, Figure 2, and page 1134, column 2 paragraphs 1-3.	14 ----- 15-16

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/20319

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 7-8, 11 and 16  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
Please See Extra Sheet.
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☒ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/20319

## A. CLASSIFICATION OF SUBJECT MATTER: IPC (7):

C07K 1/00, 14/00, 17/00; C12N 9/00, 9/12; C12P 21/06; C30B 7/00, 21/02, 28/06; G01N 31/00, 33/00, 33/53, 33/567

## BOX I. OBSERVATIONS WHERE CLAIMS WERE FOUND UNSEARCHABLE

2. Where no meaningful search could be carried out, specifically:

Claims 7-8, 11 and 16 are unsearchable to the extent that they require reference to the specified sequences from the sequence listing. Because Applicant has not included these sequences on the submitted machine-readable copy of the sequence listing, no meaningful search of the sequences per se can be carried out by this Authority. However, the subject matter of the claims has been searched to the extent possible with reference to the balance of the description.

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

Group I, claim(s) 1-12, drawn to an isolated polynucleotide, a polypeptide, and a method of assaying compound interaction with the polypeptide.

Group II, claim 13, drawn to method of preparing proteins for crystallization.

Group III, claim(s) 14-16, drawn to a process for drug design.

The inventions listed as Groups I-III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The products/processes of Groups I-III are separate and patentably distinct inventions in that the polynucleotide, polypeptide and method of compound assay of Group I, the method of preparing proteins for crystallization of Group II, and the process of drug design in Group III do not require each other for their practice and have separate functions all of which constitute the special technical features which define the contribution of each invention. Since these special technical features are not shared by each product, the inventions of Groups I-III do not form a single inventive concept within the meaning of Rule 13.2.